



K24P 4042

Reg. No. :

Name :

I Semester M.Sc. Degree (C.B.S.S. – Supplementary)

Examination, October 2024

(2021 and 2022 Admissions)

MATHEMATICS

MAT1C03 : Real Analysis

Time : 3 Hours

Max. Marks : 80

PART – A

Answer **any four** questions from this Part. **Each** question carries 4 marks.

(4×4=16)

1. Define countable set. Prove that the set of all integers is countable.
2. If p is a limit point of a set E , then prove that every neighborhood of p contains infinitely many points of E .
3. When can you say that a function f is said to be differentiable at a point x ?
Suppose f and g are defined on $[a, b]$ and are differentiable at a point $x \in [a, b]$, then prove that $f + g$ is differentiable at x and $(f + g)'(x) = f'(x) + g'(x)$.
4. Suppose f is differentiable in (a, b) . If $f'(x) = 0$ for all $x \in (a, b)$, then prove that f is constant.
5. Let f be an increasing function defined on $[a, b]$ and let x_0, x_1, \dots, x_n be $n + 1$ points such that $a = x_0 < x_1 < x_2 < \dots < x_n = b$. Then prove that $\sum_{k=1}^{n-1} [f(x_{k+1}) - f(x_k)] \leq f(b) - f(a)$.
6. Prove that $f(x) = x^2 \cos\left(\frac{1}{x}\right)$, if $x \neq 0, f(0) = 0$ is of bounded variation on $[0, 1]$.

P.T.O.



PART – B

Answer **any four** questions from this part **without** omitting **any** Unit. **Each** question carries **16** marks.

(4×16=64)

Unit– I

7. a) Let $\{E_n\}$, $n = 1, 2, 3, \dots$ be a sequence of countable sets and put $S = \cup_{n=1}^{\infty} E_n$. Then prove that S is countable.
- b) Suppose $y \subset \subset X$. A subset E of Y is open relative to Y if and only if $E = Y \cap G$ for some open subset G of X .
8. a) Show that there exist perfect sets in R^1 which contain no segment.
- b) Prove that a subset E of the real line R^1 is connected if and only if it has the following property.
If $x \in E$, $y \in E$, and $x < z < y$, then $z \in E$.
9. a) Prove that a mapping f of a metric space X into a metric space Y is continuous on X if and only if $f^{-1}(V)$ is open in X for every open set V in Y .
- b) Suppose f is a continuous mapping of a compact metric space X into a metric space Y . Then prove that $f(X)$ is compact.
- c) If f is a continuous mapping of a metric space X into a metric space Y , and if E is a connected subset of X , then prove that $f(E)$ is connected.

Unit – II

10. a) Suppose f is continuous on $[a, b]$, $f'(x)$ exists at some point $x \in [a, b]$, g is defined on an interval I which contains the range of f , and g is differentiable at the point $f(x)$. If $h(t) = g(f(t))$, ($a \leq t \leq b$), then prove that h is differentiable at x and $h'(x) = g'(f(x))f'(x)$.
- b) Give an example of a function f , which is differentiable at all points x , but f' is not a continuous function. Justify.
11. a) State and prove Taylor's theorem.
- b) Suppose f is a continuous mapping of $[a, b]$ into R^k and f is differentiable in (a, b) . Then prove that there exists $x \in (a, b)$ such that $|f(b) - f(a)| \leq (b - a)|f'(x)|$.
- c) Define Riemann-Stieltjes integral of f with respect to α over $[a, b]$.



- 12. a) Suppose f is bounded on $[a, b]$, f has only finitely many points of discontinuity on $[a, b]$, and α is continuous at every point at which f is discontinuous. Then prove that $f \in R(\alpha)$.
- b) Suppose $f \in R(\alpha)$ on $[a, b]$, $m \leq f \leq M$, ϕ is continuous on $[m, M]$ and $h(x) = \phi(f(x))$ on $[a, b]$. Then prove that $h \in R(\alpha)$ on $[a, b]$.

Unit – III

- 13. a) State and prove the formula for "integration by parts".
 - b) If f and F map $[a, b]$ into \mathbb{R}^k , if $f \in R$ on $[a, b]$, and if $F' = f$, then prove that $\int_a^b f(t)dt = F(b) - F(a)$.
 - c) If f maps $[a, b]$ into \mathbb{R}^k and if $f \in R(\alpha)$ for some monotonically increasing function α on $[a, b]$, then prove that $|f| \in R(\alpha)$ and $\left| \int_a^b f d\alpha \right| \leq \int_a^b |f| d\alpha$.
- 14. a) Let f be of bounded variation on $[a, b]$ and assume that $c \in (a, b)$. Then prove that f is of bounded variation on $[a, c]$ and on $[c, b]$ and $V_f(a, b) = V_f(a, c) + V_f(c, b)$.
 - b) Let f be of bounded variation on $[a, b]$. Let V be defined on $[a, b]$ by $V(x) = V_f(a, x)$ if $a < x \leq b$, $V(a) = 0$. Then prove that
 - i) V is an increasing function on $[a, b]$.
 - ii) $V - f$ is an increasing function on $[a, b]$.
- 15. a) Define Rectifiable paths and its arc-length. If $c \in (a, b)$ then prove that $\Lambda_f(a, b) = \Lambda_f(a, c) + \Lambda_f(c, b)$.
 - b) Consider a rectifiable path f defined $[a, b]$. If $x \in (a, b]$, let $s(x) = \Lambda_f(a, x)$ and let $s(a) = 0$. Then prove that
 - i) The function s so defined is increasing and continuous on $[a, b]$.
 - ii) If there is no subinterval of $[a, b]$ on which f is constant, then s is strictly increasing on $[a, b]$.
-