



0062687

K19U 2257

Reg. No. : .....

Name : .....

V Semester B.Sc. Degree (CBCSS- Reg./Sup./Imp.)

Examination, November-2019

(2014 Admn. Onwards)

Core Course in Mathematics

5B08 MAT: Vector Calculus

Time : 3 hrs

Max. Marks : 48

**SECTION - A**

All the 4 questions are compulsory. They carry 1 mark each. (4×1=4)

1. Find the divergence of  $e^x(\cos y \vec{i} + \sin y \vec{j})$ .
2. Express  $\frac{\partial w}{\partial r}$  in terms of  $r$  and  $s$  if  $w=x+y, x=r+s, y=r-s$ .
3. What do you mean by a potential function for a vector field  $F$ .
4. Give a parametrization of the cone  $z = \sqrt{x^2 + y^2}, 0 \leq z \leq 1$ .

**SECTION - B**

Answer any 8 questions among the questions 5 to 14. These questions carry 2 marks each. (8×2=16)

5. Find the angle between the planes  $3x-6y-2z=15$  and  $2x+y-2z=5$ .
6. Show that  $\vec{r}(t) = \cos t \vec{i} + \sqrt{5} \vec{j} + \sin t \vec{k}$  has constant length and is orthogonal to its derivative.
7. Define saddle point.

P.T.O.



8. Find the curl with respect to the right hand Cartesian coordinates of  $yz\vec{i} + 3zx\vec{j} + z\vec{k}$ .
9. Prove that for any twice continuously differentiable scalar function  $f$ ,  $\text{curl}(\text{grad } f) = \vec{0}$ .
10. Find the local extreme values of the function  $f(x,y) = xy - x^2 - y^2 - 2x - 2y + 4$ .
11. Show that  $\vec{F} = (2x - 3)\vec{i} - z\vec{j} + \cos z\vec{k}$  is not conservative.
12. Evaluate  $f(x,y,z) = 3x^2 - 2y + z$  over the line segment  $C$  joining the origin to the point  $(2,2,2)$ .
13. Find the circulation of the field  $F = (x-y)\vec{i} + x\vec{j}$  around the circle  $\vec{r}(t) = (\cos t)\vec{i} + (\sin t)\vec{j}$ ,  $0 \leq t \leq 2\pi$ .
14. Use Green's theorem to find the outward flux for the field  $F = (x-y)\vec{i} + (y-x)\vec{j}$  across the curve square bounded by  $x = 0, x = 1, y = 0, y = 1$ .

### SECTION - C

Answer any 4 questions among the questions 15 to 20. These questions carry 4 marks each. (4×4=16)

15. Find and graph the osculating circle for a parabola  $y = x^2$  at the origin.
16. Find the distance from  $S(1,1,3)$  to the plane  $3x + 2y + 6z = 6$ .
17. Find the derivative of  $f(x,y,z) = x^3 - xy^2 - z$  at  $P_0(1,1,0)$  in the direction of  $\vec{A} = 2\vec{i} - 3\vec{j} + 6\vec{k}$ . Find the direction in which  $f$  increases most rapidly at  $P$ .
18. Use Taylor's formula to find a quadratic approximation of  $f(x,y) = \cos x \cos y$  at the origin. Estimate the error in the approximation if  $|x| \leq 0.1$  and  $|y| \leq 0.1$ .
19. Integrate  $g(x,y,z) = x + y + z$  over the surface of the cube cut from the first octant by the planes  $x = a, y = a, z = a$ .
20. Find the surface area of a sphere of radius  $a$ .



## SECTION - D

Answer any 2 questions among the questions 21 to 24. These questions carry 6 marks each. (2×6=12)

21. Find:

- Unit tangent vector  $T$ ,
- Unit normal vector  $N$ ,
- Curvature  $K$ ,
- Torsion  $\tau$  and binomial vector  $B$  for the space curve  
 $\vec{r}(t) = (3\sin t)\vec{i} + 3(\cos t)\vec{j} + 4t\vec{k}$ .

22. Find the absolute maximum and minimum values of  $f(x, y) = 2 + 2x + 2y - x^2 - y^2$  on the triangular plate bounded by the lines  $x = 0, y = 0, y = 9 - x$ .

23. a) State both forms of Green's theorem.

- b) Verify the circulation -curl form of Green's theorem for the field  $\vec{F}(x, y) = (x - y)\vec{i} + x\vec{j}$  and the region  $R$  bounded by the unit circle .

$$C: \vec{r}(t) = \cos t \vec{i} + \sin t \vec{j}, 0 \leq t \leq 2\pi$$

24. a) State Stoke's theorem.

- b) Use Stoke's theorem to evaluate  $\int_C \vec{F} \cdot d\vec{r}$ , if  $F = xz\vec{i} + xy\vec{j} + 3xz\vec{k}$   $C$  is the boundary of the portion of the plane  $2x + y + z = 2$  in the first octant, traversed counter clock wise.
-