



0151443

K19P 1189

Reg. No. : .....

Name : .....

III Semester M.Sc. Degree (CBSS-Reg./Suppl./Imp.)

Examination, October - 2019

(2017 Admn. Onwards)

MATHEMATICS

MAT3C14 : ADVANCED REAL ANALYSIS

Time : 3 Hours

Max. Marks : 80

## PART - A

Answer **Four** questions from this part. Each question carries 4 marks. (4×4=16)

1. If  $\{f_n\}$  and  $\{g_n\}$  are sequences of bounded functions and converge uniformly on a set  $E$ , prove that  $\{f_n g_n\}$  converges uniformly on  $E$ .
2. Consider  $f(x) = \sum_{n=1}^{\infty} \frac{1}{1+n^2x}$ . Is  $f$  continuous wherever the series converges?
3. Show that  $e^x$  defined on  $\mathbb{R}^1$  satisfy the relation  $e^{x+y} = e^x e^y$ .
4. Show that the functional equation  $\Gamma(x+1) = x\Gamma(x)$  holds if  $0 < x < \infty$ .
5. Prove that  $\lim_{n \rightarrow \infty} \left(1 + \frac{x}{n}\right)^n = e^x$ .
6. If  $A \in L(\mathbb{R}^n, \mathbb{R}^m)$  and  $B \in L(\mathbb{R}^m, \mathbb{R}^k)$ , then prove that  $\|BA\| \leq \|B\| \|A\|$ .

## PART - B

Answer **Four** questions from this part without omitting any unit. Each question carries **16** marks. (4×16=64)

## UNIT - I

7. a) Suppose  $\lim_{n \rightarrow \infty} f_n(x) = f(x)$  ( $x \in E$ ) and put  $M_n = \sup_{x \in E} |f_n(x) - f(x)|$ .

P.T.O.



Show that  $f_n \rightarrow f$  uniformly on  $E$  if and only if  $M_n \rightarrow 0$  as  $n \rightarrow \infty$ .

- b) Suppose  $f_n \rightarrow f$  uniformly on a set  $E$  in a metric space. Let  $x$  be a limit point of  $E$ , and suppose that  $\lim_{t \rightarrow x} f_n(t) = A_n$  ( $n=1,2,\dots$ ). Then prove that  $\{A_n\}$  converges and  $\lim_{t \rightarrow x} f(t) = \lim_{n \rightarrow \infty} A_n$ .
8. a) If  $X$  is a metric space,  $C(X)$  denote the set of all complex valued, continuous, bounded functions with domain  $X$ . Show that  $C(X)$  with supremum norm is a metric space.
- b) Prove that there exists a real continuous function on the real line which is nowhere differentiable.
9. a) If  $K$  is a compact metric space, if  $f_n \in C(K)$  For  $n=1,2,\dots$  and if  $\{f_n\}$  is pointwise bounded and equicontinuous on  $K$  then prove that
- $\{f_n\}$  is uniformly bounded on  $K$ .
  - $\{f_n\}$  contains a uniformly convergent subsequence.
- b) Define equicontinuity and give an example.

## UNIT - II

10. a) Suppose the series  $\sum_{n=0}^{\infty} C_n x^n$  converges for  $|x| < R$  and define
- $$f(x) = \sum_{n=0}^{\infty} C_n x^n \quad (|x| < R).$$

Then prove that  $\sum_{n=0}^{\infty} C_n x^n$  converges uniformly on  $[-R + \epsilon, R - \epsilon]$ , no matter which  $\epsilon > 0$  is chosen. Also shows that the function  $f$  is continuous and differentiable in  $(-R, R)$ , and

$$f'(x) = \sum_{n=1}^{\infty} n C_n x^{n-1} \quad (|x| < R).$$

- b) Given a double sequence  $\{a_{ij}\}$ ,  $i=1,2,\dots$   $j=1,2,\dots$  suppose that  $\sum_{j=1}^{\infty} |a_{ij}| = b_i$  ( $i=1,2,\dots$ ) and  $\sum b_i$  converges. Then show that
- $$\sum_{i=1}^{\infty} \sum_{j=1}^{\infty} a_{ij} = \sum_{j=1}^{\infty} \sum_{i=1}^{\infty} a_{ij}.$$





11. a) Suppose  $a_0, \dots, a_n$  are complex numbers  $n \geq 1, a_n \neq 0, P(z) = \sum_{k=0}^n a_k z^k$ . Then prove that  $P(z) = 0$  for some complex number  $z$ .
- b) If, for some  $x$ , there are constants  $\delta > 0$  and  $M < \infty$  such that  $|f(x+t) - f(x)| \leq M|t|$  for all  $t \in (-\delta, \delta)$ , then prove that  $\lim_{N \rightarrow \infty} S_N(f; x) = f(x)$ .
12. a) If  $f$  is continuous (with period  $2\pi$ ) and if  $\varepsilon > 0$ , then prove there is a trigonometric polynomial  $P$  such that  $|P(x) - f(x)| < \varepsilon$  for all real  $x$ .
- b) If  $f$  is a positive function on  $(0, \infty)$  such that
- $f(x+1) = x f(x)$ .
  - $f(1) = 1$
  - $\log f$  is convex
- Then prove that  $f(x) = \Gamma(x)$ .

### UNIT - III

13. a) Suppose  $X$  is a vector space, and  $\dim X = n$ . Show that
- a set  $E$  of  $n$  vectors in  $X$  spans  $X$  if and only if  $E$  is independent
  - $X$  has a basis, and every basis consists of  $n$  vectors.
  - If  $1 \leq r \leq n$  and  $\{y_1, y_2, \dots, y_r\}$  is an independent set in  $X$ , then show that  $X$  has a basis containing  $\{y_1, y_2, \dots, y_r\}$ .
- b) Define linear transformation and give an example.
14. a) Let  $\Omega$  be the set of all invertible linear operators on  $\mathbb{R}^n$ . If  $A \in \Omega, B \in L(\mathbb{R}^n)$ , and  $\|B - A\| \cdot \|A^{-1}\| < 1$ , then prove that  $B \in \Omega$ .
- b) Suppose  $E$  is an open set in  $\mathbb{R}^n, f$  maps  $E$  into  $\mathbb{R}^m, f$  is differentiable at  $x_0 \in E, g$  maps an open set containing  $f(E)$  into  $\mathbb{R}^k$ , and  $g$  is differentiable at  $f(x_0)$ . Then prove that the mapping  $F$  of  $E$  into  $\mathbb{R}^k$  defined by  $F(x) = g(f(x))$  is differentiable at  $x_0$  and  $F'(x_0) = g'(f(x_0))f'(x_0)$ .



15. a) Suppose  $f$  maps a convex open set  $E \subset \mathbb{R}^n$  into  $\mathbb{R}^m$ ,  $f$  is differentiable in  $E$  and  $f'(x) = 0$  for all  $x \in E$ , then prove that  $f$  is constant.
- b) Suppose  $f$  maps an open set  $E \subset \mathbb{R}^n$  into  $\mathbb{R}^m$ . Then prove that  $f \in C^1(E)$  if and only if the partial derivatives  $D_j f_i$  exist and are continuous on  $E$  for  $1 \leq i \leq m, 1 \leq j \leq n$ .